



Comment on "Protein Sequences from Mastodon and Tyrannosaurus rex Revealed by Mass Spectrometry"

Mike Buckley, *et al.*
Science **319**, 33c (2008);
DOI: 10.1126/science.1147046

The following resources related to this article are available online at www.sciencemag.org (this information is current as of January 9, 2008):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/319/5859/33c>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/319/5859/33c/DC1>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/cgi/content/full/319/5859/33c#related-content>

This article **cites 12 articles**, 6 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/319/5859/33c#otherarticles>

This article appears in the following **subject collections**:

Paleontology

<http://www.sciencemag.org/cgi/collection/paleo>

Technical Comments

http://www.sciencemag.org/cgi/collection/tech_comment

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

Comment on “Protein Sequences from Mastodon and *Tyrannosaurus rex* Revealed by Mass Spectrometry”

Mike Buckley,¹ Angela Walker,² Simon Y. W. Ho,³ Yue Yang,¹ Colin Smith,⁴ Peter Ashton,¹ Jane Thomas Oates,¹ Enrico Cappellini,¹ Hannah Koon,¹ Kirsty Penkman,¹ Ben Elsworth,¹ Dave Ashford,¹ Caroline Solazzo,¹ Phillip Andrews,² John Strahler,² Beth Shapiro,⁶ Peggy Ostrom,⁵ Hasand Gandhi,⁵ Webb Miller,⁶ Brian Raney,⁷ Maria Ines Zylber,⁸ M. Thomas P. Gilbert,⁹ Richard V. Prigodich,¹⁰ Michael Ryan,¹¹ Kenneth F. Rijdsdijk,¹² Anwar Janoo,¹³ Matthew J. Collins^{1*}

We used authentication tests developed for ancient DNA to evaluate claims by Asara *et al.* (Reports, 13 April 2007, p. 280) of collagen peptide sequences recovered from mastodon and *Tyrannosaurus rex* fossils. Although the mastodon samples pass these tests, absence of amino acid composition data, lack of evidence for peptide deamidation, and association of $\alpha 1(I)$ collagen sequences with amphibians rather than birds suggest that *T. rex* does not.

Early reports of DNA preservation in multimillion-year-old bones (i.e., from dinosaurs) have been largely dismissed (1, 2) (table S1), but reports of protein recovery are persistent [see (3) for review]. Most of these studies used secondary methods of detection, but Asara *et al.* (2) recently reported the direct identification of protein sequences, arguably the gold standard for molecular palaeontology, from fossil bones of an extinct mastodon and *Tyrannosaurus rex*. After initial optimism generated by reports of dinosaur DNA, there has been increasing awareness of the problems and pitfalls that bedevil analysis of ancient samples (1), leading to a series of recommendations for future analysis (1, 4). As yet, there are no equivalent standards for fossil protein, so here we apply the recommended tests for DNA (4) to the

authentication of the reported mastodon and *T. rex* protein sequences (2) (Table 1).

First, the likelihood of collagen survival needs to be considered. The extremely hierar-

chical structure of collagen results in unusual, catastrophic degradation (5) as a consequence of fibril collapse. The rate of collagen degradation in bone is slow because the mineral “locks” the components of the matrix together, preventing helical expansion, which is a prerequisite of fibril collapse (6). The packing that stabilizes collagen fibrils (6) also increases the temperature sensitivity of degradation (E_a 173 kJ mol⁻¹) (Fig. 1). Collagen decomposition would be much faster in the *T. rex* buried in the then-megathermal (>20°C) (7) environment of the Hell Creek formation [collagen half-life ($T_{1/2}$) = ~2 thousand years (ky) than it would have been in the mastodon lying within the Doeden Gravel Beds (present-day mean temperature, 7.5°C; collagen $T_{1/2}$ = 130 ky) (Fig. 1).

This risk of contamination also needs to be evaluated. Collagen is an ideal molecular target for this assessment because the protein has a highly characteristic motif that is also sufficiently variable to enable meaningful comparison between distant taxa if enough sequence is obtained (Fig. 2). Compared with ancient DNA amplification, contamination by collagen is inherently less likely. Furthermore, because the bones sampled in (2) were excavated by the

Table 1. Key questions to ask about ancient biomolecular investigations [adapted from (4)].

Test	Sample	Pass	Observation
Do the age, environmental history, and preservation of the sample suggest collagen survival?	Mastodon, 300 to 600 ky old	Yes	Collagen $T_{1/2}$ at 7.5°C = 130 ky
	<i>T. rex</i> , 65 million years old	No	Collagen $T_{1/2}$ at 20°C = 2 ky
Do the biomolecular and/or macromolecular preservation of the sample, the molecular target, the innate nature of the sample, and its handling history suggest that contamination is a risk?	Biomolecular preservation	?	Range of evidence presented (8) but no amino acid compositional data
	Macromolecular preservation	Yes	Macromolecular preservation is not the equivalent of biomolecular preservation (9)
Do the data suggest that the sequence is authentic, rather than the result of damage and contamination?	Molecular target	Yes	Large (2.5 g) samples increase risk of contamination? Errors in interpretation of spectra [see table S1 and (13)]? Damage-induced errors in sequence
	Handling history	Yes?	
Do the results make sense, and are there enough data to make the study useful and/or to support the conclusions?	Mastodon and <i>T. rex</i>	No	Weak affinity to mammals
	Mastodon	Yes	
	<i>T. rex</i>	No	Affinity of $\alpha 1(I)$ peptides to amphibians, not birds or reptiles

¹BioArch, Departments of Biology, Archaeology, Chemistry and Technology Facility, University of York, Post Office Box 373, York YO10 5YW, UK. ²Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, MI 48109-0404, USA. ³Evolutionary Biology Group, Department of Zoology, University of Oxford, OX1 3PS, UK. ⁴Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D-04103, Leipzig, Germany. ⁵Department of Zoology, Michigan State University, East Lansing, MI 48824, USA. ⁶Department of Biology, Pennsylvania State University, University Park, PA 16802, USA. ⁷Center for Biomolecular Science and Engineering, University of California–Santa Cruz, CA 95064, USA. ⁸Department of Parasitology, Kuvim Center, Hebrew University of Jerusalem, Israel. ⁹Biological Institute, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark. ¹⁰Chemistry Department, Trinity College, 300 Summit Street, Hartford, CT 06106, USA. ¹¹Cleveland Museum of Natural History, 1 Wade Oval Drive, University Circle, Cleveland, OH 44106, USA. ¹²National Museum of Natural History “Naturalis,” P.O. Box 9517, 2300 RA Leiden, Netherlands. ¹³National Heritage Trust Fund Mauritius, Mauritius Institute, La Chaussée Street Port Louis, Mauritius.

*To whom correspondence should be addressed. E-mail: mc80@york.ac.uk




Fig. 1. Plot of radiocarbon age versus estimated effective collagen degradation temperature for radiocarbon-dated bones from laboratory databases (principally Oxford and Groningen). The line represents the expected calendar age at which 1% of the original collagen remains following a zero-order reaction; almost no bone collagen survives beyond this predicted limit. (Inset) The 99% confidence intervals of amino acid compositions by first two principal component analyses (48% of total variance) for bones from NW Europe aged <11 ky ($n = 324$), 11 to 110 ky ($n = 210$), 110 to 130 ky ($n = 26$), and 130 to 700 ky ($n = 31$). Pliocene samples are not plotted, as their composition ($n = 8$) is highly variable and yields of amino acids are low. The orange line indicates a compositional trend observed when compact bone is heated for 32 days at 95°C, which reduces collagen to 1% of the initial concentration [each inflection represents a separate analysis; $n = 32$]. The composition becomes more similar to mixed tissue samples (meat and bone meal; $n = 32$), principally due to the depletion of Gly. An amino acid profile for mammoth is consistent with collagen, unlike the associated sediment sample [data from (11)].

authors, obvious contamination sources such as animal glue (used in conservation) can be excluded. However, concentrating protein from the large amounts of bone used (2.5 g) may have heightened the risk of extraneous proteins entering the sample during extraction, although there have been no systematic studies of this phenomenon. Independent extraction and analyses would have strengthened claims for the authenticity of the origin of the peptides (and potentially ameliorated the original problems of data interpretation) (4).

The remarkable soft-tissue preservation of the investigated *T. rex* specimen (MOR 1125) has been documented (8). However, microscopic preservation does not equate with molecular preservation (9). Immunohistochemistry provides support for collagen preservation, but Asara *et al.* (2) presented no data regarding inhibition assays with collagen from different species or cross-reactivity with likely contaminants [e.g., fungi (10)]. Curiously, no amino acid compositional analysis was conducted [see (11)], although ammonium ions were identified by time-of-flight secondary ion mass spectrometry. In our experience, collagen-like amino acid profiles have been obtained in all bones from which we could obtain collagen sequence (Fig. 1, inset).

Regarding the proof of sequence authenticity, the spectra reported by Asara *et al.* (12) are inconsistent with some of the sequence assignments (13) (table S1). A common diagenetic modification, deamidation, not considered in (2), may shed light on authenticity. The facile succinimide-mediated deamidation (14) of asparagine occurred at N₂₂₉G and N₁₁₅₆G in ostrich peptides (Ost 4 and Ost5) (see table S1 for nomenclature), presumably during sample preparation. Direct hydrolytic deamidation is slower (14), and an expectation of elevated levels of such products is reasonable for old samples. We agree with the most recent interpretation (13) of the spectrum illustrated in Fig. 2B as $\alpha 1(I)$ G₃₆₂SEGPEGVR₃₇₀, the deamidated (Q→E₃₆₇) form of the sequence found in most mammals (12). By way of contrast, none of the three glutamine residues in the reported *T. rex* peptides are deamidated (table S1). Only time will tell if Q→E is a useful marker for authentically old collagen, but from the evidence presented, the mastodon sequence looks more diagenetically altered than *T. rex*.

The unusual, fragmented nature of the reported *T. rex* sequence does not make it amenable to standard, model-based phylogenetic analysis. Instead, we examined the phylogenetic

signal of the $\alpha 1(I)$ fragments of mastodon and *T. rex* using Neighbor-Net analysis and uncorrected genetic distances. Using the sequences reported in (13), both the *T. rex* and mastodon signal display an affinity with amphibians (Fig. 2A). Our reinterpretation of the spectra (12) changes the affinity of mastodon but not of *T. rex* (Fig. 2B). In addition to the $\alpha 1(I)$ peptides used in the Neighbor-Net analysis, Asara *et al.* reported two other peptides from *T. rex* (13); we question the interpretation of the $\alpha 1(II)$ spectra (identical to frog) but not the $\alpha 2(I)$ spectra (identical to chicken).

We require more data to be convinced of the authenticity of the *T. rex* collagen sequences reported by Asara *et al.* Nevertheless, the handful of spectra reported for the temperate Pleistocene mastodon fail neither phylogenetic nor diagenetic tests, thus highlighting the potential of protein mass spectrometry to bridge the present gap in our understanding between the fate of archaeological and fossil proteins. To avoid past mistakes of ancient DNA research (1), we recommend that future fossil protein claims be considered in light of tests for authenticity such as those presented here.

Reference and Notes

1. E. Willerslev, A. Cooper, *Proc. R. Soc. London. B. Biol. Sci.* **272**, 3 (2005).
2. J. M. Asara, M. H. Schweitzer, L. M. Freemark, M. Phillips, L. C. Cantley, *Science* **316**, 280 (2007).
3. M. H. Schweitzer, *Palaeontologia Electronica* **5**, editorial 2 (2003); http://palaeo-electronica.org/2002_2/r_and_p.pdf
4. M. T. P. Gilbert, H.-J. Bandelt, M. Hofreiter, I. Barnes, *Trends Ecol. Evol.* **20**, 541 (2005).
5. M. J. Collins, M. S. Riley, A. M. Child, G. Turner-Walker, *J. Archaeol. Sci.* **22**, 175 (1995).
6. C. A. Miles, M. Ghelashvili, *Biophys. J.* **76**, 3243 (1999).
7. K. R. Johnson, D. J. Nichols, J. H. Hartman, *The Hell Creek Formation and the Cretaceous-Tertiary Boundary in the Northern Great Plains: Geological Society of America Special Paper* **361**, 503–510 (2002).
8. M. H. Schweitzer *et al.*, *Science* **316**, 277 (2007).
9. N. S. Gupta, D. E. G. Briggs, R. D. Pancost, *J. Geol. Soc. London* **163**, 897 (2006).
10. P. Sepulveda *et al.*, *Infect. Immun.* **63**, 2173 (1995).




Fig. 2. Phylogenetic networks of $\alpha 1(I)$ sequences using Neighbor-Net analysis (**A**) with the most recent Asara *et al.* assignments (13) and (**B**) after our reinterpretation of the mass spectrometric data (12). *T. rex* does not group with bird/reptile using either set of sequence alignments. More sequence is required for a full, model-based phylogenetic analysis.

11. M. Schweitzer, C. L. Hill, J. M. Asara, W. S. Lane, S. H. Pincus, *J. Mol. Evol.* **55**, 696 (2002).
12. See supporting material on Science Online.
13. J. M. Asara *et al.*, *Science* **317**, 1324 (2007).
14. N. E. Robinson *et al.*, *J. Pept. Res.* **63**, 426 (2004).
15. This work was supported by NSF (EAR-0309467), National Environment Research Council (NE511148,

GR9/01656), National Center for Research Resources (P41-18627), European Commission (MEST-CT-2004-007909, MEST-CT-2005-020601), Wellcome Trust Bioarchaeology Fellowships (K.P. and H.K.) Analytical Chemistry Trust Fund, the Royal Society of Chemistry Analytical Division, Engineering and Physical Sciences Research Council, and the Michigan Proteome Consortium.

Supporting Online Material

www.sciencemag.org/cgi/content/full/319/5859/33c/DC1
SOM Text
Table S1
References

26 June 2007; accepted 20 November 2007
10.1126/science.1147046